

Patient Name SAMPLE

Patient Date of Birth dd/mm/yyyy

Test Analysis SAMPLE

Date Completed

Disclaimer

Please note we do not provide medical advice or services. If you have health disorders, medical conditions, or any condition needing medical supervision you should consult your doctor or medical professional. All products and services are provided for educational purposes and research purposes only and are not intended to be a substitute for a proper medical consultation; and the site, services, products and materials may support the relationship between you and your healthcare provider, but are not intended to replace it. They should not be used as a substitute for professional diagnosis and treatment. If you suffer from any health condition you must consult your doctor or medical professional. We do not recommend self-diagnosis or self-medication, and no information within our site or presented by us or our associates may be construed or interpreted as recommending self-diagnosis or self-medication.



PATIENT FIRST NAME :

PATIENT SURNAME:

DATE OF BIRTH:

GENDER:

ADDRESS:

Kidney Function & Liver Function Screen

BLOOD - SERUM	Result	Range	Units	
UEC (Renal)				
SODIUM	136	135 - 145	mmol/L	•
POTASSIUM	5.4	3.5 - 5.5	mmol/L	•
CHLORIDE	97	95 - 108	mmol/L	•
BICARBONATE	25	21 - 32	mmol/L	•
UREA	3.6	2.8 - 7.6	mmol/L	•
CREATININE (mmol/L)	0.06	0.04 - 0.10	mmol/L	•
Creatinine	60	40 - 100	umol/L	•
Estimated GFR	88	> 60	ml/min	
UEC Comment	eGFR : >/	= 60 mL/min/1.	73 sq.m - Do	es not exclude mild renal impairement, or kidney

eGFR : >= 60 mL/min/1.73 sq.m - Does not exclude mild renal impairement, or kidney diseases without renal impairement.

LIVER FUNCTION TESTS

BILIRUBIN (TOTAL)	14	2 - 20	umol/L
ALP	52	45 - 125	units/L
GGT	46 **H	5 - 39	units/L
ALT	48	10 - 50	units/L
AST	30	10 - 50	units/L
PROTEIN - TOTAL	73	60 - 83	g/L
ALBUMIN	42	35 - 50	g/L



LFT Comment

GAMMA-GLUTAMYLTRANSFERASE COMMENT:

GGT elevation is sensitive, but non-specific, for hepatobilary diseases. Biliary obstruction can be intrahepatic, extrahepatic or infiltrative. Microsomal GGT can also be induced by many drugs (phenytoin etc.) and ETOH, this change is reversable.

(*) Result outside normal reference range (**) Result is critically abnormal